

On the vorticity of the Universe

D. PALLE

Zavod za teorijsku fiziku

Institut Rugjer Bošković

Pošt. Pret. 180, HR-10002 Zagreb, CROATIA

Summary. — Recent analyses of the first-year WMAP data claim large-scale asymmetry and anisotropy of the CMBR fluctuations. We argue that the covariant and gauge invariant treatment of density fluctuations formulated by Ellis and Bruni can explain the asymmetric and anisotropic WMAP data by including the vorticity of the Universe. It appears that the spatial gradients of the density contrast are proportional to the vorticity of the Universe, thus allowing measurements and quantifications of the magnitude and axis of the possible cosmological rotation.

PACS 98.80 – Cosmology.

The first-year WMAP data represent the first step towards higher-precision cosmology. As expected, the WMAP data confirmed some standard cosmological settings concerning baryonic and nonbaryonic matter, primordial fluctuations, cold dark matter paradigm or spatial flatness. We should also stress unexpected findings,

such as low large-scale power of fluctuations that cannot be fitted by flat Friedmann cosmologies with a positive cosmological constant inferred from small-scale structure data, very small bound on tensor-mode fluctuations (contrary to the expectations based on the inflationary models), running of the power index at small scales and rather large reionization redshift and optical depth [1]. It is interesting to note that the integrated Sachs-Wolfe effect of the models with the negative cosmological constant reduces the large scale power of fluctuations.

Eventually, we should be forced to change both the physical content [2] and the geometry of the Universe [3] to fully understand a detailed picture growing from more refined cosmic data.

The small amount of acceleration in the geometry beyond that of Robertson-Walker can substantially change the evolution of the density contrast at small redshifts [4]. For the same initial contrast at high redshift, Eq.(13) of Ref.[4] gives the same deviation from $\Omega_m = 1$ at zero redshift for the following cosmic parameters: $\delta(\Omega_m = 1, \Sigma = 0) - \delta(\Omega_m = 2, \Sigma = 0) \simeq \delta(\Omega_m = 1, \Sigma = 0) - \delta(\Omega_m = 1, \Sigma = 3 \times 10^{-3})$, where $\Omega_m + \Omega_\Lambda = 1$ is assumed and Σ is the acceleration parameter $A_c A^c = \Sigma H^2$, $A_c = \dot{u}_c$ [4]. From such calculations one can conclude on the importance of studying more general geometries with the inclusion of acceleration and vorticity of the Universe. It is intriguing that cosmic acceleration induces anomalous constant acceleration towards the Sun [5], that is much smaller than the Pioneer anomaly $a_\Sigma = \sqrt{\Sigma} c H_0 \ll a(Pioneer)$. However, the Pioneer anomaly is too large to be consistent with planetary orbits.

Heavy Majorana neutrinos emerging from nonanomalous and nonsingular the-

ory [2] are good candidates for cold-dark-matter particles to solve the problem of the cosmic photon diffuse background [6], large optical depth observed by WMAP or super GZK cut-off cosmic rays [7]. There is also the possibility that the galactic center point source emission with no variability measured by Whipple [8] and Cangaroo II [9] could be interpreted in terms of the CDM galactic annihilation of very heavy particles with large annihilation cross sections [10]. Future measurements of Cangaroo III, Magic, H.E.S.S. and Veritas atmospheric Čerenkov telescopes will resolve this possible indirect evidence for the CDM.

Recent analyses of non-Gaussianity of WMAP data by wavelet techniques converge to the conclusion on positive non-Gaussian signals at large scales. The first findings of Vielva et al [11] were confirmed by Mukherjee and Wang [12]. Using the local curvature method Cabella et al [13] were able to reproduce observed non-Gaussian features. The N-point correlation functions from the WMAP data [14] also pointed to asymmetry at large scales. It was also shown that the fit of cosmological parameters, such as optical depth or the amplitude of fluctuations at two hemispheres, gave completely different values [15]. The most convincing analysis was performed using directional continuous spherical wavelets resulting in the 6σ detection of non-Gaussianity [16].

It is very difficult to find any physical source of such asymmetric WMAP data. We try to understand it assuming the cosmic geometry with vorticity and acceleration of the Universe. Let the metric have the form of Obukhov and Korotky [17]:

$$(1) \quad ds^2 = -dt^2 + R^2(t)(dx^2 + (1 - \Sigma)a^2(x)dy^2) \\ + R^2(t)dz^2 + 2R(t)b(x)dydt,$$

$$a(x) = e^{mx}, \quad b(x) = \sqrt{\Sigma}a(x),$$

$$A_b \equiv u^c \nabla_c u_b = b \dot{R} \delta_b^2, \quad \omega_{ab} = R b' \delta_{[a}^1 \delta_{b]}^2, \quad [ab] \equiv \frac{1}{2}(ab - ba).$$

Ellis and Bruni formulated [18] a covariant and gauge-invariant formalism for density fluctuations in inhomogeneous and nonisotropic spacetimes. Careful reanalysis [19] found an extra term with the spacelike gradient of vorticity in the evolution equation for the density contrast (Eq.(26) of Ref. [19]):

$$(2) \quad \ddot{\mathcal{D}}_{\perp a} + \mathcal{A}(t)\dot{\mathcal{D}}_{\perp a} - \mathcal{B}(t)\mathcal{D}_a + \mathcal{L}\mathcal{D}_a - \mathcal{C}(t)^{(3)}\nabla^b \omega_{ab} = 0,$$

$$\mathcal{D}_a \equiv R \frac{{}^{(3)}\nabla_a \rho}{\rho}, \quad {}^{(3)}\nabla_a f = h_a^b \nabla_b f,$$

$$h_{ab} = g_{ab} + u_a u_b, \quad u^a u_a = -1,$$

for further definitions see Ref.[19].

This additional term vanishes for the cosmic model of Obukhov and Korotky (Eq.(1)), thus we have to try another approach. It is possible to make the following local decomposition of the spatial variation of the density contrast [19]:

$$(3) \quad R^{(3)}\nabla_b \mathcal{D}_a \equiv \Delta_{ab} = W_{ab} + \Sigma_{ab} + \frac{1}{3}\Delta h_{ab},$$

$$W_{ab} \equiv \Delta_{[ab]}, \quad \Sigma_{ab} \equiv \Delta_{(ab)} - \frac{1}{3}\Delta h_{ab},$$

$$\Sigma_{ab} = \Sigma_{(ab)}, \quad \Sigma_a^a = 0, \quad \Delta = \Delta_a^a, \quad (ab) = \frac{1}{2}(ab + ba).$$

The most interesting part is the skew-symmetric part which is proportional to vorticity

$$(4) \quad W_{ab} = -R^2(1+w)\Theta\omega_{ab},$$

$$w = p/\rho, \quad \Theta = 3\frac{\dot{R}}{R}.$$

The form of primordial density fluctuations at decoupling could be fixed as in Ref.[20] $\mathcal{D}_a \propto A_a = \dot{u}_a$. Thus we only have to follow this component in the evolution

$$(5) \quad \mathcal{D}_a = \mathcal{D}e_a, \quad e_a \parallel A_a,$$

$$\delta = \frac{\delta\rho}{\rho} \equiv \mathcal{D}, \quad e_a e^a = 1, \quad e_a u^a = 0.$$

The gauge invariant and covariant formalism assumes $\delta_{CMBR} = \delta_{CMBR}(t, x, y, z)$, thus within the pure geometric formula we can infer some relationships between the cosmological observables at large scales without solving coupled evolution equations [21]. One can evaluate the left- and right-hand sides of Eq.(4), giving the following relations ($w(CMBR) = \frac{1}{3}$, $\vec{X} = R(t)\vec{x}$, subscript 0 denotes present values):

$$(6) \quad \left(\frac{\partial\delta_{CMBR}}{\partial Z}\right)_0 = 0,$$

$$\left(\frac{\partial\delta_{CMBR}}{\partial X}\right)_0 + \frac{2\omega_0}{\sqrt{\Sigma_0}}\delta_{CMBR,0} = 8R_0H_0\omega_0,$$

$$\omega^2 = \frac{1}{2}\omega_{ab}\omega^{ab}, \quad \frac{\delta\rho_{CMBR}}{\rho_{CMBR}} = 4\frac{\delta T_{CMBR}}{T_{CMBR}},$$

$$\omega_{ab} = \frac{1}{2}h_a^c h_b^d (\nabla_d u_c - \nabla_c u_d).$$

In flat geometry, one can define the cosmic scale at present as the Hubble distance $R_0 = H_0^{-1}$. Accounting for the possible relation between expansion and vorticity as in the Einstein-Cartan cosmology [3] $\omega_0 \simeq \frac{\sqrt{3}}{2}\Sigma_0 H_0$, we obtain

$$(7) \quad H_0^{-1} \left(\frac{\partial \delta_{CMBR}}{\partial X} \right)_0 = \sqrt{3}\Sigma_0 \left(4 - \frac{\delta_{CMBR,0}}{\sqrt{\Sigma_0}} \right).$$

The spatial gradient of the density contrast (in the direction perpendicular to the acceleration vector and the axis of rotation) evidently does not vanish if the vorticity of the Universe does not vanish, and vice versa. By directional continuous spherical wavelets one can estimate the spatial gradients of the density contrast at large angles and infer information on the vorticity measured a long time ago by Birch [22] ($\omega_0 = \mathcal{O}(10^{-13} yr^{-1})$) or calculated from the angular velocity of galaxies [23].

REFERENCES

- [1] SPERGEL D. N. ET AL, *ApJ Suppl.*, **148** (2003) 175.
- [2] PALLE D., *Nuovo Cimento A*, **109** (1996) 1535.
- [3] PALLE D., *Nuovo Cimento B*, **111** (1996) 671.
- [4] PALLE D., *astro-ph/0312308*, (2003) .

- [5] PALLE D., *astro-ph/0402304*, (2005) to appear in Acta Physica Slovaca.
- [6] PALLE D., *Nuovo Cimento B*, **115** (2000) 445.
- [7] PALLE D., *Nuovo Cimento B*, **118** (2003) 747.
- [8] KOSACK K. ET AL, *astro-ph/0403422*, (2004) .
- [9] TSUCHIYA K. ET AL, *astro-ph/0403592*, (2004) .
- [10] HOOPER D. ET AL, *astro-ph/0404205*, (2004) .
- [11] VIELVA P. ET AL, *ApJ*, **609** (2004) 22.
- [12] MUKHERJEE P. AND WANG Y., *astro-ph/0402602 to appear in ApJ*, (2004) .
- [13] CABELLA P. ET AL, *astro-ph/0406026*, (2004) .
- [14] ERIKSEN H. K. ET AL, *ApJ*, **605** (2004) 14.
- [15] HANSEN F. K. ET AL, *astro-ph/0406232*, (2004) .
- [16] MCEWEN J. D. ET AL, *astro-ph/0406604*, (2004) .
- [17] OBUKHOV YU. N. AND KOROTKY V. A., *Class. Quantum Grav.*, **4** (1987) 1633.
- [18] ELLIS G. F. R. AND BRUNI M., *Phys. Rev. D*, **40** (1989) 1804.
- [19] ELLIS G. F. R., BRUNI M. AND HWANG J., *Phys. Rev. D*, **42** (1990) 1035.
- [20] PALLE D., *Nuovo Cimento B*, **114** (1999) 853.
- [21] CHALLINOR A. D. AND LASENBY A. N., *Phys. Rev. D*, **58** (1998) 023001;
 CHALLINOR A. D. AND LASENBY A. N., *ApJ*, **513** (1999) 1; GEBBIE T. AND
 ELLIS G. F. R., *Ann. Phys.*, **82** (2000) 285.
- [22] BIRCH P., *Nature*, **298** (1982) 451.
- [23] LI L.-X., *Gen. Rel. Grav.*, **30** (1998) 497.